



Original Article

Prevalence and Determinants of Anxiety Among Hospitalized Children in Nowshera

Rabia Asghar¹, Qurat Ul Ain Ali¹, Neelam Ashraf¹, Bushra Tabssam¹ and Iram Sohan¹¹Aziz Bhatti Teaching Hospital, Gujrat, Pakistan

ARTICLE INFO

Keywords:

Anxiety, Children, Hospitalization, Prevalence, State-Trait Anxiety Inventory

How to Cite:Asghar, R., Ali, Q. U. A., Ashraf, N., Tabassum, B., & Sohan, I. (2025). Prevalence and Determinants of Anxiety Among Hospitalized Children in Nowshera: Anxiety Among Hospitalized Children. *Pakistan BioMedical Journal*, 8(3), 02-06. <https://doi.org/10.54393/pbmj.v8i3.1176>***Corresponding Author:**Rabia Asghar
Aziz Bhatti Teaching Hospital, Gujrat, Pakistan
is03006245957@gmail.comReceived date: 19th November, 2024Revised date: 27th February, 2025Acceptance date: 5th March, 2025Published date: 31st March, 2025

ABSTRACT

Hospital admission is a stressful experience for children. It leads to high anxiety levels. **Objectives:** To investigate the prevalence of anxiety among hospitalized children and the causes contributing to its severity. **Methods:** 118 children fulfilling inclusion criteria were selected by convenient sampling to carry out a cross-sectional observational study in Qazi Complex, Nowshera from December 2023 to May 2024. The children were aged between 9-15 years. Data were collected using the State-Trait Anxiety Inventory for Children, a validated self-reported questionnaire. Anxiety levels were classified into mild, moderate, and severe anxiety depending upon scoring. **Results:** The results showed that prior hospitalizations, experiencing painful processes, and lengthy hospital stays considerably increased the probability of severe anxiety. Children experiencing painful procedures had an odds ratio of 6.27 ($p < 0.001$) and those with a history of previous hospitalization were 4.56 times more expected to experience severe anxiety ($p < 0.001$). The key factor was extended hospital stays, with an OR of 9.09 ($p < 0.001$). Gender and age, however, did not significantly affect anxiety levels. **Conclusions:** The results emphasize the prerequisite for targeted interventions, such as decreasing hospital stays, reducing painful procedures, and providing emotional support. Understanding these features is vital for emerging attitudes to lessen anxiety in hospitalized children, thus refining their overall treatment results and well-being.

INTRODUCTION

Hospital admission is often an upsetting experience for children, as it represents an unexperienced and possibly frightening situation that interrupts their normal practices [1]. The mental disturbance of hospitalized children has also been reported by their attending nurses [2]. Anxiety is a common emotional response in children going through medical management, and prevalence of anxiety issues in hospitalized children has been progressively acknowledged in recent years especially during the pandemics [3, 4]. The World Health Organization (WHO) reports that psychological issues in children and adolescents, mainly anxiety, are on the upsurge worldwide [4]. Hospitalization may exacerbate these disorders, as kids are parted from their family members, faced with indefinite outcomes and exposed to invasive medical processes [5]. Several Studies suggest that children are mainly susceptible to developing anxiety during hospital

visits because of their intellectual skill to understand the seriousness of their disease however inadequate handling skills [6, 7]. Several research have confirmed anxiety as a considerable problem among hospitalized children. For example, a study establishes mild to severe depression and anxiety (approximately 68% and 63%, respectively) in hospitalized children [8]. Another study proves that children going through surgery stated higher levels of preoperative anxiety, which not only affect their psychological state but also increased complaints of postoperative pain and enhanced recovery time. A latest study stressed the role of parent anxiety in provoking children's anxiety levels, with children whose parents showed high stress showing more severe symptoms [9, 10]. Furthermore, children with prolonged sickness or those needing recurrent hospital admissions are at greater risk of developing anxiety conditions due to recurrent experience

to the hospital atmosphere [11]. The incidence of anxiety in hospitalized children is worrisome not only because of its direct psychological and emotional impact but also because it can hinder medical care. Anxiety can be a source of poor compliance with medical advice, prolonged hospital stays, and confrontation to treatment, additionally confounding recovery [12]. Understanding the prevalence of anxiety in hospitalized pediatric patients is significant for healthcare workers to develop directed interventions which report both the emotional and physical needs of these patients. Further research into the specific factors contributing to anxiety and the most operational ways to decrease it is essential for improving pediatric care and outcomes.

Hospitalization is a distressing experience for children and is frequently associated with increased anxiety, which can negatively affect treatment outcomes and recovery. Although previous studies have identified factors such as painful procedures, parental stress, and repeated admissions as contributors, there is limited local evidence quantifying the magnitude of these determinants in Pakistani pediatric populations. Moreover, few studies simultaneously evaluate multiple clinical and demographic predictors using standardized tools like the State-Trait Anxiety Inventory for Children. Therefore, this study aims to determine the prevalence of anxiety among hospitalized children and to identify key contributing factors, including prior hospitalization, painful procedures, and length of hospital stay, to guide targeted interventions.

METHODS

A cross-sectional study was conducted in Qazi Complex, Nowshera. 118 children fulfilling the inclusion criteria were selected for study from December 2023 to May 2024. An approval was granted by the Ethical Committee of Nowshera College of Nursing and Health Sciences (Ref No. 065/NCIV/2023). A written and verbal consent was taken from guardians/parents of patients. Patients fulfilling inclusion criteria were included in this research. Sample size was calculated by open EPI software with margin of error 0.05 and Confidence Interval (CI) 95%. Convenience Sampling technique was used for data collection. Children between age 9 to 15 years and admitted to hospital for surgical or medical management were included in the study after taking consent from guardians. Mentally retarded children, unwilling to participate, and pediatric patients

with intellectual disabilities were excluded from the study. In order to gather the information for study, a questionnaire The State-Trait Anxiety Scales Inventory for Children (STAIC) was administered for assessing trait and state anxiety, which include all variables to extract definite information. The reliability of STAIC has been proved by previous studies that indicate its internal consistency (Cronbach's alpha) in the range of 0.89 to 0.92 [13]. The anxiety levels were scored and labeled mild, moderate and severe. This questionnaire was used in the assessment of prevalence of anxiety in pediatric patients and the data were entered and analyzed using IBM-SPSS version 23.0.

RESULTS

The odds ratio (OR) of mean age group (12-15 years) is 1.43, indicating that children aged 12-15 are 43% more likely to experience severe anxiety compared to those aged 9-11 years, but this relationship is not statistically significant (p=0.351). In terms of counts, 16 out of 62 (25.8%) of the 12-15 age group experience severe anxiety, compared to 11 out of 56 (19.6%) of the 9-11 age group. Regarding gender, male have an OR of 1.19, suggesting they are 19% more likely to experience severe anxiety than female, though this is also not statistically significant (p=0.566). In numbers, 16 out of 66 (24.2%) male experience severe anxiety, compared to 11 out of 52 (21.2%) female. Another determinant, previous hospitalization, shows an OR of 4.56, meaning patients with a history of hospitalization are 4.56 times more likely to experience severe anxiety. This finding was highly significant (p<0.001). Here, 16 out of 38 (42.1%) patients with previous hospitalizations had severe anxiety, compared to 11 out of 80 (13.8%) without prior hospitalizations. Similarly, undergoing a painful procedure increases the odds by 6.27 times (p<0.001), with 18 out of 40 (45%) patients who underwent a painful procedure experiencing severe anxiety, compared to 9 out of 78 (11.5%) who did not. Finally, a hospital stay of more than 7 days shows the highest OR of 9.09, meaning these patients are 9.09 times more likely to suffer from severe anxiety (p<0.001). In terms of percentage, 19 out of 38 (50%) patients who stayed longer than 7 days experienced severe anxiety, compared to only 8 out of 80 (10%) with shorter stays. Overall, previous hospitalization, painful procedures, and extended hospital stays significantly increase the risk of severe anxiety, while age and gender show less influence (Table 1).

Table 1: Factors Influencing Anxiety among Hospitalized Patients

Characteristics	Anxiety level				Total (n=118)	p-value	OR (Severe vs Not Severe)
	Minimal or No (n=35)	Mild (n=27)	Moderate (n=29)	Severe (n=27)			
Age in Years (12-15 years)	17 (14)	11 (9)	18 (15)	16 (14)	62 (53)	0.351	1.43
Gender of Patient (Male)	20 (17)	12 (10)	18 (15)	16 (14)	66 (56)	0.566	1.19
Previous History of Hospitalization (Yes)	3 (6)	5 (4)	14 (12)	16 (14)	38 (32)	<0.001	4.56

Undergone Painful Procedure During Hospital Stays (Yes)	3 (3)	8 (7)	11 (9)	18 (15)	40 (34)	<0.001	6.27
Current Length of Stay at Hospital (>7 days)	1 (1)	3 (3)	15 (13)	19 (16)	38 (32)	<0.001	9.09

Patients with a history of hospitalization ($M=45.55$, $p<0.001$) and those undergoing painful procedures ($Mean=44.03$, $p<0.001$) showed significantly higher anxiety levels compared to those without ($Mean=25.96$ and $Mean=26.24$, respectively). Longer hospital stays (>7 days) also correlated with increased anxiety ($Mean=47.79$, $p<0.001$) versus shorter stays ($Mean=24.90$). In contrast, age (9-11 years: $Mean=29.75$, $p=0.098$; 12-15 years: $Mean=34.55$) and gender (male: $Mean=32.64$, $p=0.778$; female: $Mean=31.81$) did not significantly impact anxiety levels. Overall, the results highlight the substantial impact of hospitalization history, painful procedures, and prolonged stays on anxiety, with significant p-values (<0.001) indicating robust findings (Table 2).

Table 2: Mean Anxiety Scores by Demographic and Clinical Factors

Characteristics	N	Mean \pm SD	p-value
Age in Years			
9-11 Years	56	29.75 \pm 13.14	0.098
12-15 Years	62	34.55 \pm 17.53	
Gender of Patient			
Male	66	32.64 \pm 15.43	0.778
Female	52	31.81 \pm 16.22	
Previous History of Hospitalization			
Yes	38	45.55 \pm 14.02	<0.001
No	80	25.96 \pm 12.20	
Undergone Painful Procedure During Hospital Stay			
Yes	40	44.03 \pm 14.30	<0.001
No	78	26.24 \pm 12.78	
Current Length of Stay at Hospital			
Up-to 7 days	80	24.90 \pm 11.55	<0.001
> 7 days	38	47.79 \pm 11.49	

DISCUSSION

The results of this study demonstrate that numerous factors significantly contribute to the prevalence of anxiety among hospitalized children. Age and gender were not significant predictors of severe anxiety, whereas undergoing painful procedures, previous hospitalizations, and prolonged hospital stays were found to be critical contributors to heightened anxiety levels. These findings support and expand upon recent literature in the field. This study found no significant association between age and anxiety severity, with children aged 12-15 years only to some extent more likely to experience severe anxiety than those aged 9-11 years ($OR=1.43$, $p=0.351$). Likewise, gender was not a significant predictor ($OR=1.19$, $p=0.566$). This is steady with a previous study which also found no significant differences in anxiety levels based on gender [14, 15]. However, some literature proposes that younger children are more susceptible to anxiety due to less developed coping mechanisms [16]. The slight elevation of anxiety among older children in present study could be attributed to increased awareness of medical procedures and outcomes, but the differences were not statistically

significant. The results of current study show that children with a history of hospitalization are over four times more likely to experience severe anxiety compared to those without previous hospitalizations ($OR=4.56$, $p<0.001$). This result is in line with the work which reported that repeated hospital admissions increase anxiety levels due to anticipation of discomfort and pain [17]. The consequence of repeated experiences to the hospital atmosphere was also highlighted in the study by Zheng *et al* that stressed the psychological problem of prolonged illness and recurrent hospital stays on pediatric patients [18]. A major finding in the present study was the solid association between increased anxiety and going through painful procedures ($OR=6.27$, $p<0.001$). This result is consistent with a previous study which identified preoperative anxiety as a main distress among children going through surgery [19]. Painful procedures have been demonstrated to trigger pre-emptive anxiety, where children panic about forthcoming pain depending on past experiences. Current pain management methods can considerably drop anxiety levels and upsurge patient recovery [16]. This study underlines the need for childcare teams to reduce painful interventions when and wherever possible and arrange emotional support during required procedures. Children who stayed in the hospital for a period of more than seven days were nine times more likely to experience severe anxiety than those with shorter stay at hospital ($OR=9.09$, $p<0.001$). This strong connotation emulates findings that extended hospitalizations contribute to feelings of helplessness and separation, which aggravate anxiety indicators [20]. The current study supports the significance of decreasing hospital stay periods when medically possible and integrating interventions to decrease anxiety during prolonged admissions. It is suggested that long hospital stays, and painful procedures are the most important backers to anxiety. It emphasizes the need for targeted interventions. Recent appraisals have revealed effective involvements like parental involvement, cognitive-behavioral techniques, and play therapy in decreasing anxiety. The role of child friendly atmosphere and pain lessening approaches in anxiety management has

also been highlighted [16]. These approaches are well supported by the findings of this study, that suggest that addressing these high-risk factors could considerably improve anxiety levels. However, some of the limitations are inculcated in the study. The study results were students of Qazi Complex, Nowshera, and therefore may not be generalized to other schools that are in different zones with different socio-economic and cultural backgrounds. Also, using only the children's self-rated anxiety scores may be problematic because the children could produce biased scores by either inflating or underestimating their symptoms. However, the study has several strengths compared to previous works in the field of pediatric anxiety: First, chronic markers for severe anxiety were examined (e.g., bedtime, length of the hospital stay, presence of painful procedures) and were supported by significant statistical data. Most importantly, in contrast to numerous systematic reviews, this work establishes the size of these predictors, providing information for specific prevention strategies. Extending research to other geographical locations and different clinical communities would increase the applicability of the results and improve the understanding of the nature of anxiety in children admitted to hospitals.

However, the current study is limited by its single-center design, convenience sampling, and reliance on self-reported measures, which may introduce response bias and limit the broader applicability of the findings. Future research should involve multi-center studies with larger and more diverse samples to improve generalizability and explore additional psychosocial factors such as parental anxiety and coping strategies. Longitudinal designs are recommended to assess causal relationships and long-term psychological outcomes. Interventional studies focusing on pain management, child-friendly environments, and psychological support are also needed.

CONCLUSIONS

In conclusion, this research offers valuable consideration into the causes contributing to anxiety among hospitalized children. Although gender and age seem to have slight influence, painful procedures, prolonged hospital stays, and previous hospitalizations considerably escalate the risk of severe anxiety. These outcomes are consistent with recent literature, and they suggest the need for directed interventional strategies aimed at decreasing hospital related stressors. Strategies such as minimizing painful interventions, decreasing hospital stays, and providing psychological support can play an important role in improving the mental health and overall results of pediatric patients.

Authors' Contribution

Conceptualization: RA

Methodology: RA, BT, IS

Formal analysis: RA, QUAA, NA

Writing and Drafting: RA, QUAA, NA

Review and Editing: RA, QUAA, NA, BT, IS

All authors approved the final manuscript and take responsibility for the integrity of the work.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Daughtrey HR, Ruiz MO, Felix N, Saynina O, Sanders LM, Anand KJS. Incidence of Mental Health Conditions Following Pediatric Hospital Admissions: Analysis of a National Database. *Frontiers in Pediatrics*. 2024 Feb;12. doi:10.3389/fped.2024.1344870.
- [2] Penwill NY, Wong CT, Taylor DB, Freyleue SD, Bordogna AL, Bode RS, et al. Hospitalists' Perceptions of Pediatric Mental Health Boarding: Quality of Care and Moral Distress. *Hospital Pediatrics*. 2023 Mar;13(3):233-45. doi:10.1542/hpeds.2022-006913.
- [3] Correale C, Tondo I, Falamesca C, Capitello TG, Vigevano F, Specchio N, et al. Depression and Anxiety in Hospitalized Children with Epilepsy During COVID-19 Pandemic: Preliminary Findings of a Cross-Sectional Study. *European Psychiatry*. 2021 Apr;64(S1):670-671. doi:10.1192/j.eurpsy.2021.1780.
- [4] WHO. Mental Health of Adolescents. 2021. [Last Cited: 7th Jan 2025]. Available at: www.who.int/news-room/fact-sheets/detail/adolescent-mental-health.
- [5] Ginanjar MR, Ardianty S, Apriliyani K. Factors Related Anxiety Level On Hospitalized Children. *Masker Medika*. 2021 Oct;9(1):359-64. doi:10.52523/MASKERMEDIKA.V9I1.439.
- [6] Liang Y, Huang W, Hu X, Jiang M, Liu T, Yue H, et al. Preoperative Anxiety in Children Aged 2-7 Years Old: A Cross-Sectional Analysis of The Associated Risk Factors. *Translational Pediatrics*. 2021 Aug;10(8):2024. doi:10.21037/tp-21-215.
- [7] Delvecchio E, Salcuni S, Lis A, Germani A, Di Riso D. Hospitalized Children: Anxiety, Coping Strategies, and Pretend Play. *Frontiers in Public Health*. 2019 Sep;7:1-8. doi:10.3389/fpubh.2019.00250.
- [8] Correale C, Falamesca C, Tondo I, Borgi M, Cirulli F, Truglio M, et al. Depressive Anxiety Symptoms in

- Hospitalized Children With Chronic Illness During the First Italian COVID-19 Lockdown. *Children*.2022Aug; 9(8):1156. doi:10.3390/children9081156.
- [9] Kaynak H and Çövençer Ç. The Effect of Parents on The Hospital Anxiety Level of Hospitalized Children. *Journal Of Education And Research In Nursing*.2020 Dec; 17(4): 312-316. doi: 10.5222/head.2020.65471.
- [10] Lestari AW, Lee CK, Hayati H. The Impact of Parental Psychological Distress on Child Behavior Issues in Hospitalized Children. *La Pediatria Medica E Chirurgica*.2023Jun;45(1).doi:10.4081/pmc.2023.311.
- [11] Kadan G. Children in Hospital. Research on Prenatal, Postnatal, and Early Childhood Development.2020; 98-116). doi: 10.4018/978-1-7998-2952-2.ch006.
- [12] Çelikol Ş, Tural Büyük E, Yıldızlar O. Children's Pain, Fear, and Anxiety During Invasive Procedures. *Nursing Science Quarterly*.2019Jul;32(3):226-32. doi:10.1177/0894318419845391.
- [13] Essau CA, Muris P, Ederer EM. Reliability and Validity of the Spence Children's Anxiety Scale and the Screen for Child Anxiety-Related Emotional Disorders in German Children. *Journal of Behavior Therapy and Experimental Psychiatry*.2002Mar;33(1) :1-8. doi:10.1016/S0005-7916(02)00005-8.
- [14] Doupnik S, Hill D, Palakshappa D, Worsley D, Bae H, Shaik A, et al. Parent Coping Support Interventions During Acute Pediatric Hospitalizations. *Pediatrics*. 2017; 140(3): 1-16. doi:10.1542/peds.2016-4171.
- [15] Eijlers R, Staals L, Legerstee J, Berghmans J, Strabbing E, Van Der Schroeff M, et al. Predicting Intense Levels of Child Anxiety During Anesthesia Induction at Hospital Arrival. *Journal of Clinical Psychology in Medical Settings*.2020;28:313-322. doi :10.1007/s10880-020-09716-6.
- [16] Ahmadipour M, Sattari H, Nejad MA. Incidence and Risk Factors Related to Anxiety of Children and Adolescents Before Elective Surgery. *European Journal of Translational Myology*.2022;32(2).doi:10.4081/ejtm.2022.10449.
- [17] Liu W, Xu R, Jia J, Shen Y, Li W, Bo L. Research Progress on Risk Factors of Preoperative Anxiety in Children: A Scoping Review. *International Journal of Environmental Research and Public Health*.2022; 19(16). doi:10.3390/ijerph19169828.
- [18] Zheng K, Abraham C, Bruzzese JM, Smaldone A. Longitudinal Relationships Between Depression and Chronic Illness in Adolescents: An Integrative Review. *Journal of Pediatric Health Care*.2020;34(4): 333-45. doi:10.1016/j.pedhc.2020.01.008.
- [19] Morin H, Worsley D, Zhang X, Faerber J, Pettit AR, Kenyon CC, et al. Depression and Anxiety Symptoms During and After Pediatric Asthma Hospitalization. *Hospital Pediatrics*.2021;11(11):1272-9.doi:10.1542/hpeds.2020-000950.
- [20] Raj R, Nguyen M, Pozzo AM, Marsac ML, Vselvoshakaya O, Meadows AL. Effects of Trauma and Anxiety on Adherence in Pediatric Type 1 Diabetes. *Diabetes Spectrum*.2022;35(2):171-8.doi :10.2337/ds21-0024.