



Original Article



Parental Perceptions of Sensory Avoidance and Sleep Disturbances among Autistic Children in Lahore: A Cross-sectional Study

Eman Nasim¹, Mariam Liaquat¹, Sana Toqueer¹, Muskan Jaffar¹ and Taimoor Ahmad^{1*}¹University Institute of Physical Therapy, The University of Lahore, Lahore, Pakistan

ARTICLE INFO

Keywords:

Sensory Processing, Sleep Habits, Pediatric Sleep Problems, Sensory Avoidance Behaviours in Autistic Children

How to Cite:Nasim, E., Liaquat, M., Toqueer, S., Jaffar, M., & Ahmad, T. (2025). Parental Perceptions of Sensory Avoidance and Sleep Disturbances among Autistic Children in Lahore: A Cross-sectional Study: Parental Perceptions of Sensory Avoidance and Sleep Disturbances among Autistic Children. *Pakistan BioMedical Journal*, 8(6), 21-26. <https://doi.org/10.54393/pbmj.v8i6.1258>***Corresponding Author:**Taimoor Ahmad
University Institute of Physical Therapy, The University of Lahore, Lahore, Pakistan
taimoor121214@gmail.comReceived Date: 3rd May, 2025Revised Date: 19th June, 2025Acceptance Date: 21st June, 2025Published Date: 30th June, 2025

ABSTRACT

As literature shows, children with autism often experience sensory processing difficulties and sleep disturbances. **Objectives:** To determine parent-reported sensory avoidance behaviours and sleep habits in school-aged children with autism spectrum disorder in Lahore. **Methods:** This observational cross-sectional study was conducted on 249 autistic children. Outcome measures of this study were the prevalence of sensory avoidance behaviours and sleep habits. The standardized tools were the Short Sensory Profile (SSP) and the Children's Sleep Habits Questionnaire (CSHQ) to assess the sensory avoidance behaviours and sleep problems. Data were gathered by using SSP and CSHQ. Data were analyzed by using SPSS-27. **Results:** 38% were aged 3-7 years, and 61% were aged 8-12 years. Parent reports showed that 17% of children exhibited probable differences and 53% exhibited significant differences in tactile sensory avoidance behaviours. Movement sensitivity issues were definite in 54% and probable in 30% of children. Regarding auditory sensory avoidance, 55% showed significant changes, and 1% showed probable changes. For under-responsive behaviour. For visual and auditory sensory behaviours, only 44% showed typical performance. Overall, 74% of children demonstrated definite sensory avoidance behaviours across multiple domains, while 8% showed some level of sensory avoidance. Additionally, 20% did not display low energy behaviours. Regarding sleep habits, 16% had typical sleep patterns, 3% faced sleep issues at a minimum level, 22% faced sleep issues at an intermediate level, and the remaining 57% faced sleep issues at a severe level. A p-value was >0.05. Significant association between sensory avoidance behaviour and sleep habits. **Conclusion:** Sensory avoidance behaviour had a significant association with sleep habits.

INTRODUCTION

Autism Spectrum Disorder (ASD) is a complex neurodevelopmental condition characterized by deficits in social interaction, communication, and repetitive behaviours [1]. Globally, its prevalence has risen significantly over the past two decades, although regional estimates vary. The global prevalence of autism spectrum disorder (ASD) remained uncertain, with studies reporting significant variations across different regions. In low-income countries, underdiagnosis was prevalent due to limited access to assessment tools. Autism, originally coined to describe the self-isolating tendencies in individuals with schizophrenia, was later redefined in the

1940s by psychiatrists Leo Kanner and Hans Asperger. They characterized it as a distinct syndrome in children, marked by atypical social interaction, communication challenges, and repetitive behaviours. In contemporary discourse, the term Autism encompasses a diverse variety of neurological issues, unified by shared clinical features such as speech impairments, social difficulties, and restricted or repetitive behavioural patterns [1, 2]. Autism, along with other neurodevelopmental disorders, had been recognized as possessing a substantial genetic foundation; however, it had also been posited that environmental influences contributed meaningfully to its aetiology. Gene expression



in autism has varied due to copy number variations and exposure to environmental toxins [3-5]. Some cases of autism have been linked to spontaneous genetic mutations [6]. These mutations had affected neuron movement, axon direction, and the formation of synapses. Structural genetic changes were found more frequently in autistic children than in typically developing peers [7, 8]. A specific gene mutation, CHD8, had been associated with autism and was linked to physical traits such as an enlarged head and widely spaced eyes [3]. Autism Spectrum Disorder (ASD) usually develops in early life, where the issue is diagnosed as early as three years old. The initial symptoms included such signs as a poor reaction to the name and uneasiness with eye contact [9-11]. Though most social and cognitive challenges were carried down into adulthood, communication skills may prove to change with time, especially during adolescent years. Intelligence and IQ are known to be rather fixed. Nevertheless, the quality of life may improve greatly with the help of effective social support systems, interventions, and accommodations [12, 13]. Data from 2020 indicated that Hong Kong, South Korea, the United States, Japan, and Ireland had the highest recorded rates, with Hong Kong reporting 372 cases per 10,000 children [5]. The growing incidence of autism spectrum disorder (ASD) could have been explained by the fact that there are increasing environmental risk factors that are likely to increase during the prenatal period, including exposure to air pollution, advanced age of a father, and maternal use of psychotropic drugs. Though there has been a critical influence of genetic predisposition, which influences ASD, environmental factors are believed to be major contributors to the development of ASD. They may be causing the increase of ASD if the environmental factors emerged more and more strongly with the progress of time [6]. Difficulties in sensory processing are also common to a great proportion of children with ASD [14]. Sensory processing in the clinical practice was the increased sensitivity to external stimuli, sounds, lights or odors [15]. The idea of sensory integration was pioneered and explained how people perceived, processed and used rudiments of sense [16]. Researcher contribution is the basis of therapeutic approaches to working on sensory input/awareness, self-management, motor performance, and praxis. Winnie Dunn devised this model in 1997 and divided the responses into the model of sensory processing; this was dependent on the neurological threshold and behavioural response. There are neurological thresholds: lower (having a high degree of reactivity to low stimulation) and generally higher. Depending on the behavioural response, thresholds work against or with one another, resulting in four sensory

processing patterns being low registration, sensory seeking, sensory sensitivity, and sensory avoiding [7]. Sensory integration dysfunction (SID) is a disorder of communication, social interaction that is built on incorrect behavioural patterns. It elevates due to poor neural transmission between the sensory receptors, afferent systems, and other subcortical components, causing inadequacies in the reception of those stimuli, processing, and modulation of the stimulus. SID occurred as hypersensitivity, with even low stimuli provoking the inappropriate responses or hyposensitivity, requiring high stimuli to arouse neural responses. The malfunction traverses in tactile, vestibular, proprioceptive, auditory, visual, gustatory and olfactory systems that affect motor coordination, emotional steadiness and mental development [8]. Sleep disturbance is yet another of the commonly co-morbid challenges in autistic children. Instead, it has been found that sleep problems, including bedtime resistance, delayed sleep onset, and night awakenings, are very common in ASD, with more than 50% [9].

Autism Spectrum Disorder is frequently associated with sensory processing difficulties and significant sleep disturbances that adversely affect child functioning and family well-being. Although international evidence highlights a relationship between sensory avoidance behaviours and poor sleep outcomes, limited region-specific data from Lahore, Pakistan, examine this association based on parental perceptions using standardized tools. Therefore, this study aimed to determine the prevalence of sensory avoidance behaviours and sleep disturbances, and to examine their association among school-aged autistic children in Lahore. This study aims to investigate parental perceptions of sensory avoidance and sleep habits among school-aged children with autism spectrum disorder (ASD) in Lahore. Disrupted sleep exacerbated the cognitive decline in these children, further hindering their developmental progress. The findings provided valuable insights for occupational physiotherapists in formulating targeted sensory integration strategies to mitigate sensory avoidance behaviours. Implementing such interventions not only enhanced the well-being of autistic children but also served as a crucial alleviating factor for parental distress.

METHODS

This study followed a cross-sectional, descriptive design with correlational analysis and was conducted over a period of four months (January 2025 to Apr 2025) in private autistic centers located in Lahore, Pakistan. When a prevalence (p) of 73% was assumed (based on the previous

literature), with a confidence level (Z) of 95% (Z=1.96) and a margin of error (d) of 5% (d=0.05), the sample size approximately equaled 303. Based on practical limitations and scarcity of resources, a sample size of 249 participants was recruited, which was considered satisfactory to address the descriptive and correlational aims of the study [17]. In this study, a selected sample of 249 participants from the total population was used for the collection of data from private autistic centers of Lahore. The study followed the ethical guidelines and ethical approval was taken. Non-probability sampling technique was used for this purpose. The outcomes of this study were assessing sensory avoidance and sleep habits using standardized tools: The Short Sensory Profile (SSP) and the Children's Sleep Habits Questionnaire (CSHQ). Both are validated tools commonly used in neurodevelopmental research. To assess the internal consistency of the tools in the present sample, Cronbach's alpha was calculated. The SSP showed a Cronbach's alpha of 0.89, while the CSHQ yielded a Cronbach's alpha of 0.83, indicating good reliability in this population. Children aged 3–12 years with Autism Spectrum Disorder (ASD), diagnosed according to DSM-5 criteria by a licensed clinician, were included in the study. Primary caregivers were involved in the child's care and provided detailed reports of the child's sensory and sleep behaviours. Children were excluded if they had a primary diagnosis of another developmental disorder, such as intellectual disability or a genetic syndrome, which might affect sensory or sleep patterns. Children on sedative medications, with severe medical or neurological conditions, or with co-existing psychiatric disorders were excluded. Single parents with more than one disabled child, or parents not residing in the same household as the child, were also excluded. Data were analyzed using SPSS version 27.0. Categorical variables were described as absolute frequencies (n) and relative frequencies (%), while continuous variables were expressed as mean ± standard deviation for parametric distributions or as median and percentiles for non-parametric distributions. The Chi-square test was used to assess associations between categorical variables. Pearson's or Spearman's correlation coefficients were applied, as appropriate, to examine relationships between continuous variables. A p-value of less than 0.05 was considered statistically significant.

RESULTS

This study comprised 249 autistic children. Out of them, 96 autistic children (38.5%) were in the age range of 3-7 years old, and 153 autistic children (61.4%) belonged to the 8 years to 12 years' age group. The average coded age value was 1.61+/-0.49 (Table 1).

Table 1: Descriptive Statistics for Age and Gender Distribution of Autistic Children (n=249)

Variables	Descriptive Statistics
Mean	1.6145
Standard Deviation	0.48770
Minimum value	1.0
Maximum Value	2.0
Total	249

The gender ratio included 123 boys (49.40%), 126 girls (50.60%) (Figure 1).




Figure 1: Gender Distribution of Autistic Children Showing 123 Were Boys and 126 Were Girls

Sensory avoidance behaviour was assessed using the Short Sensory Profile (SSP). 41 children showed typical performance, 22 subjects had shown some issues in sensory avoidance behaviours processing, and the remaining 186 had shown significant differences in sensory avoidance behaviours processing (Table 2).

Table 2: Frequency of Sensory Avoidance Behaviours Assessed by Short Sensory Profile

Sensory Avoidance Behaviours	Frequency (%)
Typical Performance	41 (16.41%)
Probable Difference	22 (8.84%)
Significant Difference	186 (74.70%)
Total	249 (100%)

Sleep patterns were evaluated using the Children's Sleep Habits Questionnaire (CSHQ). Results showed that 42 children (16.9%) had typical sleep habits, 8 (3.2%) had mild sleep problems, 55 (22.1%) had moderate sleep problems, and 144 (57.8%) experienced severe sleep disturbances (Table 3).

Table 3: Distribution of Sleep Habits among Autistic Children

Sleep Habits	Frequency (%)
Typical Sleep Habits	42 (16.87%)
Mild Sleep Problems	8 (3.21%)

Moderate Sleep Problems	55(22.09%)
Severe Sleep Problems	144 (57.83%)
Total	249(100%)

A Chi-square test showed a significant association between age group and sensory avoidance behaviour ($\chi^2 = 9.22$, $df=2$, $p=0.010$). However, no significant association was found between gender and sensory avoidance ($\chi^2 = 0.78$, $df=2$, $p=0.675$). There was a statistically significant association between sensory avoidance behaviours and sleep disturbances ($\chi^2 = 15.62$, $df=6$, $p=0.016$), indicating that children with greater sensory avoidance had more severe sleep issues (Table 4).

Table 4: Chi-Square Test Showing Association Between Sensory Avoidance, Age, Gender, And Sleep Habits

Variable Pair	χ^2 Value	df	p-Value	Interpretation
Age Group × Sensory Avoidance	9.22	2	0.010	Significant
Gender × Sensory Avoidance	0.78	2	0.675	Not Significant
Sensory Avoidance × Sleep Habits	15.62	6	0.016	Significant

DISCUSSION

Autism Spectrum Disorder (ASD) typically manifests during early childhood, with formal diagnoses frequently established by the age of three. Early indicators often comprised reduced responsiveness to auditory cues, such as one's name being called, and a pronounced aversion to maintaining eye contact [18, 19]. Although difficulties in social interaction and cognitive functioning generally persisted into adulthood, communicative abilities often showed improvement over time, particularly during adolescence [20, 21]. Cognitive aptitude, including intelligence quotient (IQ), tended to remain stable throughout the lifespan [22]. Nevertheless, the overall quality of life for individuals with ASD was markedly enhanced through the implementation of robust social support systems, tailored therapeutic interventions, and context-specific accommodations [23]. This observational study included children between the ages of 3 and 12 years who had autism spectrum disorder. Information was collected from their parents. According to parent reports, 17% of the children showed possible differences, and 53% showed clear differences in how they avoided touch. For taste and smell, 32% had clear differences and 12% had possible differences. Problems with movement sensitivity were definite in 54% of the children and possible in 30%. When it came to avoiding sounds, 55% showed clear signs and 1% showed possible signs. For under-responsiveness (less reaction to sensory input), 48% showed definite signs and 1% showed possible signs. Only 44% of the children showed normal behaviour in both seeing and hearing responses. Overall, 74% of the children had definite sensory avoidance in more than one area, and 8% had some level of sensory avoidance. Also,

20% of the children did not show signs of low energy behaviours. Olson, investigating the school-aged children with autism, found links between one-carbon metabolism (OCM) nutrients and sensory processing patterns, revealing significant associations between intake of vitamins B1, B12, and choline with specific sensory domains [24]. In contrast, the present study focused on parent-reported sensory avoidance behaviours and sleep habits without assessing nutritional factors. While the earlier research highlighted biological influences on sensory responses, our findings emphasized the behavioural manifestations of sensory avoidance across multiple domains, along with a notable association with sleep difficulties. Both studies underscore the complexity of sensory processing in autism, though from different perspectives biochemical versus behavioural. These differences suggest a need for integrative approaches in future research to better understand the interplay between diet, sensory behaviours, and overall well-being in autistic children. Aljaid examined sleep habits among autistic children under 15 years and found significant differences compared to typically developing peers, including shorter sleep duration, earlier bedtimes, and higher rates of sleep disturbances such as snoring and morning headaches [25]. Similarly, the present study revealed that a large proportion (57%) of autistic children experienced severe sleep problems, with only 16% displaying typical sleep patterns. Gulati reported that 77.5% of children with ASD were poor sleepers, supported by objective polysomnographic findings showing reduced sleep efficiency and altered sleep architecture [26].

A key limitation of this study is its cross-sectional design and use of non-probability sampling from private centers, which limits causal inference and generalizability to the broader ASD population. Additionally, reliance on parent-reported questionnaires may introduce reporting bias. Future research should adopt longitudinal, multicenter designs with probability sampling and incorporate objective sleep assessments (e.g., actigraphy or polysomnography) alongside sensory evaluations to better understand causal pathways and inform targeted sensory integration interventions.

CONCLUSIONS

It was concluded that sensory avoidance behaviour had a significant association with sleep habits. Early sensory integration therapies are recommended as they be employed in therapy programs in order to enhance sleep among the autistic children. Furthermore, education and training programs oriented to parents should be developed to control sensory challenges at home.

Authors' Contribution

Conceptualization: EN

Methodology: ML, ST

Formal analysis: EN

Writing and Drafting: EN, MJ, TA

Review and Editing: EN, ML, ST, MJ, TA

All authors approved the final manuscript and take responsibility for the integrity of the work.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Sauer AK, Stanton J, Hans S, Grabrucker A. Autism spectrum disorders: etiology and pathology. Exon Publications. 2021 Aug; 1-5. doi: 10.36255/exonpublications.autismspectrumdisorders.2021.etiology.
- [2] Amaral DG. Examining the causes of autism. In *Cerebrum: The Dana forum on brain science*. 2017 Jan;(2017): cer-01.
- [3] Bhat S, Acharya UR, Adeli H, Bairy GM, Adeli A. Autism: cause factors, early diagnosis and therapies. *Reviews in the Neurosciences*. 2014 Dec; 25(6): 841-50. doi:10.1515/revneuro-2014-0056.
- [4] Masi A, DeMayo MM, Glozier N, Guastella AJ. An overview of autism spectrum disorder, heterogeneity and treatment options. *Neuroscience Bulletin*. 2017 Apr; 33: 183-93. doi: 10.1007/s12264-017-0100-y.
- [5] Nkonyane F, Gurayah T, Lingah T. Challenges and coping strategies of parents of children with autism spectrum disorder in Eswatini. *Journal of Child and Adolescent Mental Health*. 2021 Sep; 33(1-3): 32-41. doi: 10.2989/17280583.2023.2259428.
- [6] Taylor MJ, Rosenqvist MA, Larsson H, Gillberg C, D'Onofrio BM, Lichtenstein P *et al.* Etiology of autism spectrum disorders and autistic traits over time. *Journal of the American Medical Association Psychiatry*. 2020 Sep; 77(9): 936-43. doi: 10.1001/jamapsychiatry.2020.0680.
- [7] Van den Boogert F, Klein K, Spaan P, Sizoo B, Bouman YH, Hoogendijk WJ *et al.* Sensory processing difficulties in psychiatric disorders: A meta-analysis. *Journal of Psychiatric Research*. 2022 Jul; 151: 173-80. doi:10.1016/j.jpsychires.2022.04.020.
- [8] Ilić-Savić I, Petrović-Lazić M, Resimić R. Sensory integration and its significance for functioning and development of children's speech. *Medicinski Pregled*. 2021; 74(5-6): 205-10. doi: 10.2298/MPNS2106205I.
- [9] Mahaseth PK and Choudhary A. Sensory Integration Therapy Verses Conventional Physical Therapy Among Children with Cerebral Palsy On Gross Motor Function-A Comparative Randomized Controlled Trial. *Annals of the Romanian Society for Cell Biology*. 2021Apr; 25(4).
- [10] Kuhlthau K, Orlich F, Hall TA, Sikora D, Kovacs EA, Delahaye J *et al.* Health-related quality of life in children with autism spectrum disorders: Results from the autism treatment network. *Journal of Autism and Developmental Disorders*. 2010 Jun; 40: 721-9. doi: 10.1007/s10803-009-0921-2.
- [11] Halstead EJ, Joyce A, Sullivan E, Tywyn C, Davies K, Jones A *et al.* Sleep disturbances and patterns in children with neurodevelopmental conditions. *Frontiers in Pediatrics*. 2021 Mar; 9: 637770. doi: 10.3389/fped.2021.637770.
- [12] Draudvilienė L, Sosunkevič S, Daniusevičiūtė-Brazaitė L, Burkauskienė A, Draudvila J. The benefit assessment of the physiotherapy sessions for children with autism spectrum disorder. *Baltic Journal of Sport and Health Sciences*. 2020 Nov; 3(118): 25-32. doi: 10.33607/bjshs.v3i118.971.
- [13] Warutkar VB, Kovala RK, Warutkar V. Review of sensory integration therapy for children with cerebral palsy. *Cureus*. 2022 Oct; 14(10). doi: 10.7759/cureus.30714.
- [14] Guller B and Yaylaci F. Eating and sleep problems, related factors, and effects on the mental health of the parents in children with autism spectrum disorder. *International Journal of Developmental Disabilities*. 2024 May; 70(3): 406-15. doi: 10.1080/20473869.2022.2095689.
- [15] Griffin ZA, Boulton KA, Thapa R, DeMayo MM, Ambarchi Z, Thomas E *et al.* Atypical sensory processing features in children with autism, and their relationships with maladaptive behaviors and caregiver strain. *Autism Research*. 2022 Jun; 15(6): 1120-9. doi: 10.1002/aur.2700.
- [16] Hologue C, Volk H, Crocetti D, Gottlieb B, Spira AP, Mostofsky SH. Links between parent-reported measures of poor sleep and executive function in childhood autism and attention deficit hyperactivity disorder. *Sleep Health*. 2021 Jun; 7(3): 375-83. doi: 10.1016/j.sleh.2020.12.006.
- [17] Ben-Sasson A, Hen L, Fluss R, Cermak SA, Engel-Yeger B, Gal E. A meta-analysis of sensory modulation symptoms in individuals with autism spectrum disorders. *Journal of Autism and Developmental Disorders*. 2009 Jan; 39: 1-1. doi: 10.1007/s10803-008-0593-3.

- [18] Pickard H, Hirsch C, Simonoff E, Happé F. Exploring the cognitive, emotional and sensory correlates of social anxiety in autistic and neurotypical adolescents. *Journal of Child Psychology and Psychiatry*. 2020 Dec; 61(12): 1317-27. doi: 0.1111/jcpp.13214.
- [19] Horiuchi F, Kawabe K, Oka Y, Nakachi K, Hosokawa R, Ueno SI. The association between autistic traits and sleep habits/problems in toddlers. *Developmental Neuropsychology*. 2020 Dec; 45(7-8): 485-95. doi: 10.1080/87565641.2020.1865357.
- [20] Glod M, Riby DM, Rodgers J. Relationships between sensory processing, repetitive behaviors, anxiety, and intolerance of uncertainty in autism spectrum disorder and Williams syndrome. *Autism Research*. 2019 May; 12(5): 759-65. doi: 10.1002/aur.2096.
- [21] Kadwa RA, Sahu JK, Singhi P, Malhi P, Mittal BR. Prevalence and characteristics of sensory processing abnormalities and its correlation with FDG-PET findings in children with autism. *The Indian Journal of Pediatrics*. 2019 Nov; 86: 1036-42. doi: 10.1007/s12098-019-03061-9.
- [22] Burns CO, Dixon DR, Novack M, Granpeesheh D. A systematic review of assessments for sensory processing abnormalities in autism spectrum disorder. *Review Journal of Autism and Developmental Disorders*. 2017 Sep; 4: 209-24. doi: 10.1007/s40489-017-0109-1.
- [23] Al-Farsi OA, Al-Farsi YM, Al-Sharbaty MM, Al-Adawi S. Sleep habits and sleep disorders among children with autism spectrum disorders, intellectual disabilities and typically developing children in Oman: a case-control study. *Early Child Development and Care*. 2019 Dec; 189(14): 2370-80. doi: 10.1080/03004430.2018.1455671.
- [24] Olson A, Krall JR, Baranova A, Slavin M. Nutritional Intake and Sensory Processing in School-Aged Children with Autism Spectrum Disorder. *Nutrients*. 2025 Feb; 17(4): 604. doi: 10.3390/nu17040604.
- [25] Aljaid MS, Alharthi MS, Alrubie SS, Qattan FM, Almalki AS, Alzahrani ES et al. Comparative Analysis of Sleep Behavior in ASD Children: A Taif City Perspective. *Bangladesh Journal of Medical Science*. 2025 Feb: 123-32. doi: 10.3329/bjms.v24i10.79181.
- [26] Gulati S, Sharma S, Singh A, Chowdhary SR, Meena A, Singh S et al. 1167 The spectrum of Co-morbidities in children with ASD (Autism Spectrum Disorder). *Archives of Disease in Childhood*. 2022 Aug; 107(Suppl 2): A323-4. doi: 10.1136/archdischild-2022-rcpch.523.