

PAKISTAN BIOMEDICAL JOURNAL

https://www.pakistanbmj.com/journal/index.php/pbmj/index ISSN(E): 2709-2798, (P): 2709-278X Volume 8, Issue 11 (November 2025)



Original Article



Quality of Life among Women with Pelvic Organ Prolapses After Postpartum

Aneeqa Aqdas¹, Mahnoor Alam¹, Nimra Rehman¹, Areeba Ahsan Qadri¹, Rimsha Tariq¹, Fatima Mazhar¹, Erum Ghaffar¹, Maria Samad¹, Uzma Irshad¹ and Barjees Ahmad¹

¹Faculty of Allied Health Sciences, Hajvery University, Lahore, Pakistan

ARTICLE INFO

Keywords:

Pelvic Organ Prolapse, Quality of Life, Postpartum Women, P-OoL Questionnaire

How to Cite:

Aqdas, A., Alam, M., Rehman, N., Qadri, A. A., Tariq, R., Mazhar, F., Ghaffar, E., Samad, M., Irshad, U., & Ahmad, B. (2025). Quality of Life among Women with Pelvic Organ Prolapses After Postpartum: Women with Pelvic Organ Prolapses After Postpartum. Pakistan BioMedical Journal, 8(11), 28–32. https://doi.org/10.54393/pbmj.v8i11.1292

*Corresponding Author:

Aneeqa Aqdas Faculty of Allied Health Sciences, Hajvery University, Lahore, Pakistan aneeqach99@gmail.com

Received Date: 24th September, 2025 Revised Date: 17th November, 2025 Acceptance Date: 23rd November, 2025 Published Date: 30th November, 2025

ABSTRACT

Pelvic organ prolapse (POP) is the dropping of the pelvic organs as a result of the inability to hold them through weakness, especially in the aftermath of childbirth, which causes physical and emotional problems. **Objectives:** To evaluate the prevalence of POP and its effects on the quality of life (QoL) of postpartum women by administering the P-QoL questionnaire. Purposes: To establish the QoL of postpartum women who have pelvic organ prolapse. **Methods:** The study was a three-month cross-sectional study to gather information on various hospitals within Lahore. In the present study, 377 postpartum women (20 to 45 years old) in the public hospitals were evaluated by the P-QoL questionnaire, but they were not included whose prior POP treatment or were still pregnant. **Results:** The researchers established that 96.3 percent of 377 women experienced severe symptoms of POP, with the moderate-to-high performance scores consistent, and only half of them made it to the standardized analysis. **Conclusions:** The fact that severe POP symptoms occur in the majority of cases (96.3%), and moderate-to-high performance consistency highlights the fact that there is a strong necessity to enhance data inclusion and specific interventions.

INTRODUCTION

Pelvic Organ Prolapse (POP) is a common problem, as weakened pelvic support results in dropper organs such as the bladder or uterus, which can cause the effect of bladder, bowel, and sexual problems [1, 2]. It is largely associated with giving birth, aging, obesity, and hard work. POP has a strong impact on the quality of life of women, yet it is usually underdiagnosed because of its stigma and inconsistent symptoms, as well as lack of adequate care access, particularly in low-resource environments. Physiotherapy or pessaries can be used in the management of cases at an early stage [3]. The knowledge of the POP effects and especially among the postpartum women, will play a crucial role in quick intervention and subsequent

improvement in health [4]. The situation is further aggravated by the fact that the condition is more common in the low-income areas with restricted access to healthcare services, including gynecological surgeries [5]. As much as the current body of research progressively increases on the prevalence of POP and its effects on the quality of life, little is known regarding the effects that the condition has on the postpartum period, a period that is already fraught with a lot of emotional and physical transformations in women [6, 7]. Symptoms of prolapse at early stages, often beginning after delivery, have not been given much attention, and the literature is predominantly dedicated to general female groups or to the cases of the

latter stage of prolapse [8, 9]. Moreover, the limited studies of postpartum POP that are presently available are more likely to underestimate its manifold effects on daily functioning, psychology, and physical recovery. Moreover, the perspectives of physiotherapy are not fully included, particularly in the assessment of the impact of dysfunction of the pelvic floor on postpartum quality of life. Therefore, the postpartum population remains underrepresented in care paradigms and clinical research [10].

This study aimed to explore the association between the quality of life influenced by pelvic organ prolapses following childbirth and the significance of enhanced postpartum screening and management interventions.

METHODS

The cross-sectional research design was used to identify the prevalence and related variables of pelvic organ prolapse at a given time. This design, however, was restrictive by nature in developing causal relationships between identified risk factors and the condition. The data was gathered by various hospitals in Lahore. This would be a three-month (July 2025 to September 2025) study period, which included ethical clearance of the Hajvery University with ref no: HU/FAHS/ECRB/2461, recruitment of the study participants, and data collection through the use of the P-QoL questionnaire, data analysis, and compiling of the report. A simple random sampling strategy was undertaken so that all eligible people in the sampling frame got an equal opportunity of being selected. The study sample comprised all multiparous postpartum women (20-45, up to 6 months after delivery) attending the gynecology Outpatient Departments of the study hospitals in Lahore within the four months of the study. Potential participants were assigned sequential numbers, and a computer-based random number generator was used to select the required 377 participants from this complete list. The sample size was calculated using the Raosoft Sample Size Calculator. For the calculation, an expected prevalence of pelvic organ prolapses of 50% was used, as no definitive local data was available, and this proportion maximizes the required sample size for a given margin of error. With a 95% confidence level and a 5% margin of error, the calculated minimum sample size was 377 participants. The target population was women aged 20-45 after postpartum. Inclusion criteria included women aged 20-45 years, who had delivered within the postpartum period (up to 6 months after childbirth), multiparous (having had two or more deliveries) women were included, as repeated childbirth is a known risk factor for pelvic organ prolapse, and women with the ability to understand and complete the P-QoL questionnaire (self-administered or interview-assisted). Exclusion criteria were primiparous women (first delivery) to ensure focus on multiparity-related risk factors, women with ongoing pregnancy or multifetal gestation, women with a history of prior pelvic organ prolapse treatment, pelvic floor reconstructive surgery, or hysterectomy, women diagnosed with neurological, systemic, or cognitive impairments that could interfere with questionnaire completion, women with a history of perinatal death, chronic illnesses, or mental health disorders and women currently undergoing pelvic floor physiotherapy or conservative management for prolapse. Eligible participants were informed about the study and asked to sign an informed consent form. P-QoL questionnaire was given either as a self-administered or an interview using Urdu (translated when necessary). The Prolapse Quality of Life (P-QoL) Questionnaire was a validated instrument that was employed to determine the effect of pelvic organ prolapse. A strict forward and backward translation was carried out into the Urdu language to be used in this study. First, forward translation was done by two bilinguals who are independent. The translated versions were then harmonized into one draft by a committee of health care professionals and linguists. This was then back-translated to English by two other translators who were not told what the original instrument was. The back-translated one was put against the original to verify conceptual, semantic, and content equivalency. The last Urdu version was pilot-tested by a group of 20 postpartum women (not included in the main study), to ensure that there were comprehension and cultural appropriateness, and understanding. The analysis of data was based on version SPSS version 23.0. Demographic and clinical variables were summarized using descriptive statistics (mean, standard deviation, frequencies, and percentages). The prevalence was determined as the percentage of women who had POP symptoms. Inferential statistics was used to enhance the analytical scope to go beyond description. The Chi-square test was applied to determine the connection between categorical variables, including age groups, and the severity of POP impact on QoL. The p-value was taken to be significant at less than 0.05.

RESULTS

The number of women enrolled in the study was 377, who developed pelvic organ prolapse (POP). All the 50 participants excluded in the final analysis were those whose responses to the questionnaire were incomplete, and therefore, 327 valid cases were used to conduct the statistical analysis (86.7%). During clinical examination, the severity of POP was determined using the standardized system of Pelvic Organ Prolapse Quantification (POP-Q). This system marks prolapse centimeters of the descent of the hymen relative to the pelvic organs between 0 and 4. To make this analysis, some stages were grouped as follows: Stage 0-II was categorized as 'Moderate', and Stage III-IV

was categorized as 'Severe (Table 1).

Table 1: Severity Level of POP

Severity Level	Frequency (%)
Moderate	14 (3.7%)
Severe	363 (96.3%)
Total	377 (100.0%)

The relationship between the age group and the extent to which POP affects the QoL was evaluated statistically. A Chi-square test of independence showed that there was a significant relationship between these variables, χ^2 (12, n=377) = 45.8, p<0.001. The highest percentage (31.6%) of women who were severely affected were in the age group 3640 years old, stating they experienced the highest impact(Table 2).

Table 2: Age and Severity of Respondents Affecting QoL

Age Group (Years)	Frequency (%)	Severity of Impact
20-25	50 (13.3%)	Mildly Affected
26-30	70 (18.5%)	A Lot Affected
31–35	80 (21.3%)	Moderately Affected
36-40	118 (31.6%)	Severely Affected
41-45	59 (15.3%)	Little Affected
Total	377 (100.0%)	_

Levels of symptom distress were uneven, as the results of the Pelvic Organ P-QOL questionnaire were found. The top complaints were bowel issues (10.3%), sexual discomfort (9.0%), and lower back pain with vaginal pain (8.2%). Common urinary symptoms included poor urinary stream (5.6%), straining to empty the bladder (5.8%), and urine dribbling (4.5%). Psychosocial impacts such as depression (1.1%), sleep disturbance (1.1%), and restricted social life (0.8%) were reported less frequently (Table 3).


Table 3: Distribution of P-QOL Domain Scores

Variables	Symptom/ Domain, n (%)
Bowels Do Not Feel Empty	39 (10.3%)
Vaginal Bulge Interfering with Sex	34 (9.0%)
Lower Backache with Vaginal Discomfort	31(8.2%)
Straining to Open Bowels	27(7.2%)
Constipation	26(6.9%)
Straining To Empty Bladder	22 (5.8%)
Poor Urinary Stream	21(5.6%)
Urine Dribbles After Emptying	17 (4.5%)
Other Minor Symptoms	<4% Each
Total	377 (100.0%)

Out of 377 participants, 118 women (31.6%) reported being severely affected, 80(21.3%) were a lot affected, 70(18.5%) were moderately affected, and 59 (15.3%) were a little affected. For the analysis of the overall quality of life impact, data from 50 participants (13.3%) were excluded. This exclusion was necessary because these participants

did not complete critical sections of the P-QOL questionnaire, specifically one or more domains required to compute a valid total QOL impact score (Figure 1).

Quality of Life (Total Women = 377)



■ Excluded ■ Severely Affected ■ Moderately Affected □ A Little Affected

Figure 1: Quality of Life Affected by POP

DISCUSSION

The present study finds that the prevalence of pelvic organ prolapse (POP) is very high among postpartum women in Pakistan, with the majority having moderate to severe symptoms and a serious quality of life impairment, particularly in particular, physical and emotional wellbeing. The majority was middle-aged, and care-seeking was low due to stigma and lack of awareness. Compared to global data, the burden is higher in low-resource settings due to limited screening and postpartum support. While pelvic floor exercises have proven effective elsewhere, few women in this study practiced them, highlighting the need for greater physiotherapy integration and postpartum care improvements. Fourteen 3241-year-old women with parity 13 attended, POP Grade 13. The reports of women being dismissive or not being appreciative of the effect of their condition were high. Other reports involved communication with the HCPs who they believed were listening, understood the impact of their POP, positive diagnoses, and realistic treatment procedures [11]. Previous findings indicate the need to carry out larger epidemiological studies on the potential adverse reproductive health effects of heavy load-bearing exercises on third-world females [12]. Prevalence was 30.9 percent (95 percent confidence interval: 24.4-38.2 percent), (heterogeneity 12=99.8 percent). The metaanalysis of subgroups in the studies delivered the questionnaire to assess the prevalence rate and found that the prevalence was 25.0% and the physical examination

was 41.8% [13]. One of the studies demonstrated the possibility of a pessary to positively influence the quality of life of women and the opportunity to considerably improve the sexual functioning and body perception [14]. High rates of LAM avulsion were detected in Chinese women with a POP of 39%. It was associated with an elevated level of POP, and these women were more handicapped in QoL. However, LAM avulsion was not an independent issue that influenced the QoL of such women [15]. POP also affects women, and the majority of them who visit the gynecology clinic in MRRH have the disease in stage II. POP is advised to be regularly screened to enable the detection and preventive action to be taken, especially with grand multiparous, peasant farmers, and people who have a history of perineal tears [16]. This was analyzed in 1426 women. There was a 4.9% prevalence of POP. POP was significantly related to the parity [17]. High rates of prevalence of depression in women with advanced prolapse of the pelvic organs were also reported by a study in comparison to the rest of the studies. The screening measures to be employed in the detection and management of depression in women who have advanced prolapse of the pelvic organs should be formulated to help identify the condition at its early stages [18]. A shared aetiology of both POP and other disorders that are related to collagen may occur at the molecular level of the collagens [19]. The awareness of the complexity of the etiology, epidemiology, and management of POP is significant in minimizing its burden and maximizing the health and well-being of the populations affected [20].

CONCLUSIONS

This research concluded that the fact that severe POP symptoms (96.3%) were high and consistent performance moderate-high should prompt the need to improve data collection and specific interventions. A study that gauges the incidence of depression and anxiety among postpartum women with POP should be conducted. Research needs to be conducted in the future to address women lived experiences, cultural beliefs, and barriers to care using the qualitative method.

Authors Contribution

Conceptualization: AA¹ Methodology: NR, AA² Formal analysis: MS, UI, BA

Writing review and editing: AA, MA, RT, FM, EG

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Belayneh T, Gebeyehu A, Adefris M, Rortveit G, Gjerde JL, Ayele TA. Pelvic Organ Prolapse Surgery and Health-Related Quality of Life: A Follow-Up Study. BioMed Central Women's Health. 2021 Jan; 21(1): 4. doi: 10.1186/s12905-020-01146-8.
- [2] Hage-Fransen MA, Wiezer M, Otto A, Wieffer-Platvoet MS, Slotman MH, Nijhuis-Van Der Sanden MW et al. Pregnancy-and Obstetric-Related Risk Factors for Urinary Incontinence, Fecal Incontinence, or Pelvic Organ Prolapse Later in Life: A Systematic Review and Meta-Analysis. Acta Obstetricia Et Gynecologica Scandinavica. 2021 Mar; 100(3): 373-82. doi: 10.1111/aogs.14027.
- [3] Doğan K, Öztoprak MY, Dura MC, Aslan İÖ. The Effect of Stress Incontinence and Pelvic Organ Prolapse Surgery on Sexual Function and Quality of Life. Journal of the Turkish German Gynecological Association. 2024 Jun; 25(2): 96. doi: 10.4274/jtgga. galenos.2024.2023-1-13.
- [4] Dora BT, Kassa ZY, Hadra N, Tsigie BB, Esayas HL. Determinants of Pelvic Organ Prolapse at Public Hospitals in Hawassa City, Southern Ethiopia, 2020: Unmatched Case Control Study. BioMed Central Women's Health. 2022 Jul; 22(1): 301. doi: 10.1186/s 12905-022-01890-z.
- [5] Hendrix SL, Clark A, Nygaard I, Aragaki A, Barnabei V, McTiernan A. Pelvic Organ Prolapse in the Women's Health Initiative: Gravity and Gravidity. American Journal of Obstetrics and Gynecology. 2002 Jun; 186(6): 1160-6. doi: 10.1067/mob.2002.123819.
- [6] Mukwege A, Harlow SD, Hood MM, Reed BD, Dugan SA, Miller JM. Race/Ethnicity and Incidence of Pelvic Organ Prolapse in Midlife Women: The Study of Women's Health Across the Nation. Journal of Women's Health. 2024 Aug; 33(8): 1042-51. doi: 10.1089/jwh.2023.0804.
- [7] Reimers C, Stær-Jensen JE, Siafarikas F, Bø K, Engh ME. Association Between Vaginal Bulge and Anatomical Pelvic Organ Prolapse During Pregnancy and Postpartum: An Observational Study. International Urogynecology Journal. 2018 Mar; 29(3): 441-8. doi: 10.1007/s00192-017-3407-3.
- [8] Robinson D, Prodigalidad LT, Chan S, Serati M, Lozo S, Lowder J et al. International Urogynaecology Consultation Chapter 1 Committee 4: Patients'

- Perception of Disease Burden of Pelvic Organ Prolapse. International Urogynecology Journal. 2022 Feb; 33(2): 189-210. doi: 10.1007/s00192-021-04997-3.
- [9] Rebmann E, Hamel JF, Helbert C, Lemasson F, Legendre G, Venara A. Anal Incontinence After Obstetrical Anal Sphincter Injury Significantly Impacts Quality of Life for Women: A Cohort Study. Langenbeck's Archives of Surgery. 2024 Feb; 409(1): 67. doi: 10.1007/s00423-024-03257-4.
- [10] Brown HW, Hegde A, Huebner M, Neels H, Barnes HC, Marquini GV et al. International Urogynecology Consultation Chapter 1 Committee 2: Epidemiology of Pelvic Organ Prolapse: Prevalence, Incidence, Natural History, and Service Needs. International Urogynecology Journal. 2022 Feb; 33(2): 173-87. doi: 10.1007/s00192-021-05018-z.
- [11] Carroll L, Sullivan CO, Doody C, Perrotta C, Fullen BM. Pelvic Organ Prolapse: Women's Experiences of Accessing Care and Recommendations for Improvement. BioMed Central Women's Health. 2023 Dec; 23(1): 672. doi: 10.1186/s12905-023-02832-z.
- [12] Koyuncu A, Kadota JL, Mnyippembe A, Njau PF, Sijali TR, McCoy SI et al. Heavy Load Carrying and Symptoms of Pelvic Organ Prolapse among Women in Tanzania and Nepal: An Exploratory Study. International Journal of Environmental Research and Public Health. 2021 Feb; 18(3): 1279. doi: 10.3390/ijerph18031279.
- [13] Hadizadeh-Talasaz Z, Khadivzadeh T, Mohajeri T, Sadeghi M. Worldwide Prevalence of Pelvic Organ Prolapse: A Systematic Review and Meta-Analysis. Iranian Journal of Public Health. 2024 Mar; 53(3): 524. doi:10.18502/ijph.v53i3.15134.
- [14] De Albuquerque Coelho SC, De Castro EB, Juliato CR. Female Pelvic Organ Prolapse Using Pessaries: Systematic Review. International Urogynecology Journal. 2016 Dec; 27(12): 1797-803. doi: 10.1007/s 00192-016-2991-y.
- [15] Yu CH, Chan SS, Cheung RY, Chung TK. Prevalence of Levator Ani Muscle Avulsion and Effect on Quality of Life in Women with Pelvic Organ Prolapse. International Urogynecology Journal. 2018 May; 29(5): 729-33. doi: 10.1007/s00192-017-3454-9.
- [16] Tugume R, Lugobe HM, Kato PK, Kajabwangu R, Kanyesigye H, Masembe S et al. Pelvic Organ Prolapse and Its Associated Factors among Women Attending the Gynecology Outpatient Clinic at A Tertiary Hospital in Southwestern Uganda. International Journal of Women's Health. 2022 Apr: 625-33. doi: 10.2147/IJWH.S355461.

- [17] Gabra MG, Tessier KM, Fok CS, Nakib N, Oestreich MC, Fischer J. Pelvic Organ Prolapse and Anal Incontinence in Women: Screening with A Validated Epidemiology Survey. Archives of Gynecology and Obstetrics. 2022 Sep; 306(3): 779-84. doi: 10.1007/s 00404-022-06510-7.
- [18] Abebe SA, Gashaw ZM, Ayichew Z, Angaw DA, Kindie EA. Prevalence and Associated Factors of Depression among Women with Advanced Pelvic Organ Prolapse in Northwest Ethiopia: Cross-Sectional Study. BioMed Central Women's Health. 2024 May; 24(1): 313. doi: 10.1186/s12905-024-03162-4.
- [19] Lammers K, Lince SL, Spath MA, van Kempen LC, Hendriks JC, Vierhout ME et al. Pelvic Organ Prolapse and Collagen-Associated Disorders. International Urogynecology Journal. 2012 Mar; 23(3): 313-9. doi: 10.1007/s00192-011-1532-y.
- [20] Zumrutbas AE. Understanding Pelvic Organ Prolapse: A Comprehensive Review of Etiology, Epidemiology, Comorbidities, and Evaluation. Société Internationale d'Urologie Journal. 2025 Feb; 6(1): 6. doi: 10.3390/siuj6010006.